Question	Answer	Marks	Guidance
8(c)	$(AC^2) = r^2 + 0.25r^2 - (2 \times r \times 0.5r \cos(\pi - 1.45))$	**M1**	For correct substitution in cosine rule using $(\pi -$ *their* $1.45)$
	$AC = 1.17r$	**A1**	
	Perimeter = $2.95r + 1.17r$	**B1**	**FT** on *their AC*
	$r = 2.91$	**A1**	
9(a)	$\overrightarrow{AB} = \mathbf{b} - \mathbf{a}$ or $\overrightarrow{BA} = \mathbf{a} - \mathbf{b}$	**B1**	
	$\overrightarrow{OX} = \mathbf{a} + \frac{3}{4}\overrightarrow{AB}$ or $\overrightarrow{OX} = \mathbf{b} + \frac{1}{4}\overrightarrow{BA}$ $\overrightarrow{OX} = \mathbf{a} + \frac{3}{4}(\mathbf{b} - \mathbf{a})$ or $\overrightarrow{OX} = \mathbf{b} + \frac{1}{4}(\mathbf{a} - \mathbf{b})$	**M1**	For correct use of ratio, using *their* $\overrightarrow{AB}$ or $\overrightarrow{BA}$
	$\overrightarrow{OX} = \frac{\mathbf{a}}{4} + \frac{3}{4}\mathbf{b}$	**A1**	
9(b)	$\overrightarrow{AC} = 2\mathbf{b} - \mathbf{a}$	**B1**	
9(c)	$\overrightarrow{AY} = -\mathbf{a} + h\left(\frac{\mathbf{a}}{4} + \frac{3}{4}\mathbf{b}\right)$	**B1**	**FT** on *their* $\overrightarrow{OX}$
9(d)	$-\mathbf{a} + h\left(\frac{\mathbf{a}}{4} + \frac{3}{4}\mathbf{b}\right) = m(2\mathbf{b} - \mathbf{a})$	**M1**	For equating appropriate vectors and attempt to equate like vectors
	$-1 + \frac{h}{4} = -m$	**A1**	**FT** from *their* $\overrightarrow{AY}$ and $\overrightarrow{AC}$
	$\frac{3h}{4} = 2m$	**A1**	**FT** from *their* $\overrightarrow{AY}$ and $\overrightarrow{AC}$
	$h = \frac{8}{5}, \ m = \frac{3}{5}$	**A1**	For both
10(a)	$\frac{3x + 10 + 2(x+1)}{(x+1)(3x+10)} = \frac{3x + 10 + 2x + 2}{(x+1)(3x+10)}$ $= \frac{5x + 12}{3x^2 + 13x + 10}$	**B1**	For expansion and simplification to obtain given answer