Question	Answer	Marks	Guidance
9(a)	$d = 12$	**B1**	
	$\frac{n}{2}\left(-8+(n-1)12\right) > 2000$ $3n^2 - 5n - 1000 > 0$	**M1**	For use of sum formula to obtain a three term quadratic inequality or equation
	$n = \frac{5 \pm \sqrt{25+12000}}{6}$ $n = 19.1$	**M1**	**Dep** for attempt at critical value(s) using *their* quadratic, may be using a calculator, so may be implied by a correct answer of 20.
	$n = 20$	**A1**	
9(b)(i)	$r = 3$	**2**	**M1** For $ar^6 = 27$ and $ar^8 = 243$ with an attempt to eliminate a to obtain r^2. Allow other valid methods.
9(b)(ii)	3^{26}	**2**	**B1** for $a = \frac{1}{27}$ or 3^{-3} nfww
9(c)	Common ratio or $r = \sin\theta$	**B1**	May be implied by e.g. $\frac{1}{1-\sin\theta}$ or $\frac{1-\sin^n\theta}{1-\sin\theta}$
	$-1 < \sin\theta < 1$ or $\lvert\sin\theta\rvert < 1$ or $-1 < r < 1$ or $\lvert r\rvert < 1$ with no incorrect statements seen.	**B1**	**Dep** on previous B1
10(a)	$\frac{1}{\sin\alpha} + \frac{1}{\cos\alpha} \quad (=0)$	**B1**	For dealing correctly with $\text{cosec}^2\alpha$ and $\sec^2\alpha$ to obtain an expression in $\sin\alpha$ and $\cos\alpha$ only
	$\tan\alpha = -1$ or $\sin\alpha = -\cos\alpha$	**B1**	For an equation in $\tan\alpha$, may be implied by a correct solution.
	$\alpha = -\frac{\pi}{4}$ or -0.785 $\alpha = \frac{3\pi}{4}$ or 2.36	**2**	**B1** for one correct solution **B1** for a second correct solution and no extra solutions in the range.