Question	Answer	Marks	Partial Marks
4(b)(i)	$\left(1-\frac{1}{2}x^{-\frac{1}{2}}\right)\sec^2\left(x-\sqrt{x}\right)$ oe, isw	**2**	**M1** for $\mathrm{f}(x)\sec^2\left(x-\sqrt{x}\right)$
4(b)(ii)	Correctly writes $\left(1-\frac{1}{2}x^{-\frac{1}{2}}\right)\sec^2\left(x-\sqrt{x}\right)=$ $\frac{2\sqrt{x}-1}{2\sqrt{x}}\sec^2(x-\sqrt{x})$ or $\frac{2\sqrt{x}-1}{2\sqrt{x}\cos^2(x-\sqrt{x})}$ **and** states an answer $k\tan\left(x-\sqrt{x}\right)$ or states $\frac{1}{2}\int\frac{2\sqrt{x}-1}{\sqrt{x}\cos^2(x-\sqrt{x})}\,\mathrm{d}x=\tan(x-\sqrt{x})$	**M1**	where k is a non-zero constant; dependent on **part (b)(i)**
	$2\tan\left(x-\sqrt{x}\right)+c$ nfww	**A1**	
5	Correct quotient rule: $\frac{\mathrm{d}y}{\mathrm{d}x}=\frac{(\ln 3x)[1]-x\left(\frac{1}{3x}\times 3\right)}{(\ln 3x)^2}$ oe OR correct product rule using $y = x(\ln 3x)^{-1}$: $\frac{\mathrm{d}y}{\mathrm{d}x}=x\left(-(\ln 3x)^{-2}\times\frac{3}{3x}\right)+[1](\ln 3x)^{-1}$	**2**	**M1** for $\frac{\mathrm{d}y}{\mathrm{d}x}=\frac{(\ln 3x)[1]-x\times \textit{their}\left(\frac{1}{3x}\times 3\right)}{(\ln 3x)^2}$ OR for $\frac{\mathrm{d}y}{\mathrm{d}x}=x\times \textit{their}\left(-(\ln 3x)^{-2}\times\frac{3}{3x}\right)$ $+[1](\ln 3x)^{-1}$
	$\frac{\delta y}{h}=\frac{\ln 3-1}{(\ln 3)^2}$ oe, soi	**M1**	**FT** *their* $\left.\frac{\mathrm{d}y}{\mathrm{d}x}\right\vert_{x=1}$ providing quotient rule or appropriate product rule attempted
	$\delta y=\frac{\ln 3-1}{(\ln 3)^2}h$ or $\delta y = 0.0817h$ nfww	**A1**	must have evidence of correct derivative
6	$\left[-\frac{1}{4}\mathrm{e}^{2-4x}\right]_{-0.25}^{0.5}$ oe	**B2**	**B1** for $k\mathrm{e}^{2-4x}$, $k \neq -4$
	Correct use of correct limits: $-\frac{1}{4}\mathrm{e}^{0}-\left(-\frac{1}{4}\mathrm{e}^{3}\right)$ oe	**M1**	**FT** *their* $-\frac{1}{4}\mathrm{e}^{2-4x}$ providing **B1** awarded
	$-\frac{1}{4}+\frac{1}{4}\mathrm{e}^{3}$ or exact equivalent, isw	**A1**	

© Cambridge University Press & Assessment 2024